

From resolvent estimates to rates of decay

Charles Batty (University of Oxford)
with Yuri Tomilov et al

Functional Calculus and Harmonic Analysis of Semigroups
Besançon, 1 October 2014

A damped wave equation

$$\begin{aligned}\frac{\partial^2 u}{\partial t^2} - \Delta u + 2a(x) \frac{\partial u}{\partial t} &= 0 & (t > 0, x \in \Omega) \\ u(x, t) &= 0 & (t > 0, x \in \partial\Omega) \\ u(\cdot, 0) = u_0 \in H_0^1(\Omega), & \quad \frac{\partial u}{\partial t}(\cdot, 0) = u_1 \in L^2(\Omega).\end{aligned}$$

Here, Ω is a compact Riemannian manifold (with boundary), and $a : \Omega \rightarrow [0, \infty)$ (continuous).

Energy

$$E(u, t) = \frac{1}{2} \int_{\Omega} \left(|\nabla u|^2 + \left| \frac{\partial u}{\partial t} \right|^2 \right) dx,$$

decreasing in t .

Except in degenerate cases,

- 1 the energy $E(u, t) \rightarrow 0$ as $t \rightarrow \infty$;
- 2 if the domain of damping $\{x : a(x) > 0\}$ satisfies the geometric optics condition then the decay occurs at an exponential rate (Bardos-Lebeau-Rauch, 1992);
- 3 in other cases, the decay occurs at a polynomial rate or a logarithmic rate, uniformly for smooth initial data.

Reformulate the damped wave equation:

$$X = H_0^1(\Omega) \times L^2(\Omega),$$

$$A = \begin{pmatrix} 0 & 1 \\ \Delta & -2a(x) \end{pmatrix},$$

$$D(A) = (H^2 \cap H_0^1) \times H_0^1.$$

$$U(t) = \begin{pmatrix} u(t) \\ \frac{\partial u}{\partial t} \end{pmatrix} \in X, \quad E(u, t) = \frac{1}{2} \|U(t)\|_{H^1 \times L^2}^2,$$

$$U'(t) = AU(t), \quad U(0) = \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}.$$

Lebeau (1996) established that

$$\sigma(A) \subseteq \{\lambda \in \mathbb{C} : -2\|a\|_\infty \leq \operatorname{Re} \lambda < 0\},$$

and $\|(is - A)^{-1}\|$ grows (at most) polynomially as $|s| \rightarrow \infty$.

The damped wave equation is well-posed, so A generates a C_0 -semigroup of contractions $\{T(t) : t \geq 0\}$ on X .

Decay of $E(u, t)$ for initial data $(u_0, u_1) \in D(A)$ corresponds to decay of $\|T(t)(\lambda - A)^{-1}\|$ for any $\lambda \in \rho(A)$. One establishes such decay by some form of complex inversion of Laplace transforms.

An abstract theorem

X complex Banach space, $\{T(t) : t \geq 0\}$ C_0 -semigroup, generator A

If T is bounded, $\sigma(A) \subseteq \{\lambda \in \mathbb{C} : \operatorname{Re} \lambda \leq 0\}$

Theorem

If T is bounded and $\sigma(A) \cap i\mathbb{R}$ is empty, then

$$\lim_{t \rightarrow \infty} \|T(t)A^{-1}\| = 0.$$

There is also a converse result.

Consequence:

$$\lim_{t \rightarrow \infty} \|T(t)x\| = 0 \quad (x \in X).$$

$M : \mathbb{R}_+ \rightarrow (0, \infty)$, continuous, increasing.

$$M_{\log}(s) = M(s)(\log(1 + M(s)) + \log(1 + s)).$$

Theorem (M_{\log} -Theorem; B-Duyckaerts 2008)

Suppose that T is bounded and

$$\|(is - A)^{-1}\| \leq M(|s|) \quad (s \in \mathbb{R}).$$

Then, for any $c \in (0, 1)$,

$$\|T(t)A^{-1}\| = O\left(\frac{1}{M_{\log}^{-1}(ct)}\right) \quad (t \rightarrow \infty).$$

Examples

1. $M(s) = C$: one obtains exponential decay of $\|T(t)A^{-1}\|$. (True more generally, Weis-Wrobel)
2. $M(s) = C \log(2 + s) \implies \|T(t)A^{-1}\| = O\left(e^{-c\sqrt{t}}\right)$
3. $M(s) = C \exp(\alpha s)$: one obtains logarithmic decay (Lebeau, Burq for Hilbert space)
4. $M(s) = C(1 + s)^\alpha$: one obtains polynomial stability:

$$\|(is - A)^{-1}\| \leq C(1 + |s|)^\alpha \implies \|T(t)A^{-1}\| = O\left(\left(\frac{\log t}{t}\right)^{\frac{1}{\alpha}}\right)$$

slightly sharper than Liu and Rao (Hilbert space) and Batkai et al (Banach spaces)

Theorem

Let $m : (0, \infty) \rightarrow (0, \infty)$ be decreasing with $\lim_{t \rightarrow \infty} m(t) = 0$.
Assume that

$$\|T(t)(1 - A)^{-1}\| \leq m(t) \quad (t > 0).$$

Then $\sigma(A) \cap i\mathbb{R}$ is empty, and, for each $c \in (0, 1)$,

$$\|(is - A)^{-1}\| = O(m^{-1}(c/|s|)) \quad (|s| \rightarrow \infty).$$

So the apparently optimal rate of decay in Ingham-Karamata Theorem would have M^{-1} instead of M_{\log}^{-1} .

Theorem (Borichev-Tomilov 2010)

Let $M(s) = C(1 + s)^\alpha$. On general Banach spaces, it is not possible to improve the conclusion of the M_{\log} -theorem that the rate of decay is

$$O\left(\left(\frac{\log t}{t}\right)^{\frac{1}{\alpha}}\right) \quad (t \rightarrow \infty).$$

In the case of a bounded C_0 -semigroup on a Hilbert space satisfying

$$\|(is - A)^{-1}\| \leq C(1 + |s|)^\alpha \quad (s \in \mathbb{R}),$$

one has

$$\|T(t)A^{-1}\| = O\left(\frac{1}{t^{\frac{1}{\alpha}}}\right) \quad (t \rightarrow \infty).$$

Regularly varying case

M is *regularly varying* if $M(s) \sim \frac{s^\alpha}{\ell(s)}$ where ℓ is slowly varying.

Consider the case $\alpha = 1$ (purely for simplicity), ℓ increasing.

Theorem (B-Chill-Tomilov, to appear, JEMS)

Assume that X is a Hilbert space, and ℓ is slowly varying and increasing, and

$$\|(is - A)^{-1}\| = O\left(\frac{|s|}{\ell(|s|)}\right) \quad (|s| \rightarrow \infty).$$

Then

$$\|T(t)A^{-1}\| = O\left(\frac{1}{t\ell(t)}\right) \quad (t \rightarrow \infty).$$

For many (but not all) ℓ , this gives the optimal result

$$\|T(t)A^{-1}\| = O(1/M^{-1}(t)).$$

Theorem (B-Chill-Tomilov, to appear, JEMS)

Assume that X is a Hilbert space, and ℓ is slowly varying and decreasing, and

$$\|(is - A)^{-1}\| = O\left(\frac{|s|}{\ell(|s|)}\right) \quad (|s| \rightarrow \infty).$$

Then, for every $\varepsilon > 0$,

$$\|T(t)A^{-1}\| = O\left(\frac{(\log t)^\varepsilon}{t\tilde{\ell}(t)}\right) \quad (t \rightarrow \infty).$$

Here $\tilde{\ell}$ is a slowly varying function which is sometimes, but not always, the same as ℓ . However the optimal result would have $\varepsilon = 0$.

$$\Omega_M := \left\{ \lambda \in \mathbb{C} : \operatorname{Re} \lambda > -\frac{1}{M(|\operatorname{Im} \lambda|)} \right\}$$

Theorem (Ingham, Karamata, 1935; , B-Duyckaerts 2008)

Let $f \in L^\infty(\mathbb{R}_+, X)$, and assume that \hat{f} extends analytically to Ω_M and the extension satisfies

$$\|\hat{f}(\lambda)\| \leq M(|\operatorname{Im} \lambda|), \quad \lambda \in \Omega_M.$$

Let $c \in (0, 1)$. Then there exist positive numbers C and t_0 , depending only on $\|f\|_\infty$, M and c , such that

$$\left\| \hat{f}(0) - \int_0^t f(s) ds \right\| \leq \frac{C}{M_{\log}^{-1}(ct)}, \quad t \geq t_0.$$

The semigroup theorem follows by taking $f(t) = T(t)x$.

Theorem (B-Borichev-Tomilov)

Let $f \in L^p(\mathbb{R}_+, X)$, where $1 \leq p \leq \infty$, and assume that \widehat{f} extends analytically to Ω_M and satisfies

$$\|\widehat{f}(\lambda)\| \leq M(|\operatorname{Im} \lambda|), \quad \lambda \in \Omega_M.$$

Then there exists $c > 0$, depending only on p , such that the function

$$t \mapsto M_{\log}^{-1}(ct) \left(\widehat{f}(0) - \int_0^t f(s) ds \right)$$

belongs to $L^p(\mathbb{R}_+, X)$.

The shape of Ω_M is much more important than the bound on \widehat{f} .

Theorem

Let $(T(t)) : t \geq 0$ be a C_0 -semigroup on a Banach space X , with generator A , and let $\omega \in \rho(A)$, $S \in \mathcal{B}(X)$. If $x \in X$ is such that

- (i) $ST(\cdot)x \in L^p(\mathbb{R}_+, X)$ for some $p \geq 1$, and
- (ii) $SR(\cdot, A)x$ extends to an analytic function G on an open set Ω containing $\overline{\mathbb{C}_+}$,

then

$$t \mapsto ST(t)R(\omega, A)x \in L^p(\mathbb{R}_+, X).$$

If $\Omega = \Omega_M$ and, for some $\alpha, \beta > 0$,

$$\|G(\lambda)\| \leq C(1 + |\operatorname{Im} \lambda|)^\alpha M(|\operatorname{Im} \lambda|)^\beta, \quad \lambda \in \Omega_M,$$

then, for some $k > 0$,

$$t \mapsto M_{\log}^{-1}(kt)ST(t)R(\omega, A)x \in L^p(\mathbb{R}_+, X).$$

Outline of proof: M_{\log} -theorems

Newman-Korevaar approach to the Ingham-Karamata Theorem via contour integrals

Choose parameters of contour carefully as functions of t ; use pointwise estimates for various terms

$p < \infty$: Use that arc-length on the contour in the left-half plane is a Carleson measure; obtain an L^p -estimate.

Outline of proof: Hilbert space, polynomial case

Bounded semigroup on Hilbert space, $M(s) = C(1 + s)^\alpha$.

- Use complex analysis to show

$$\|(\lambda - A)^{-1}(-A)^{-\alpha}\| \leq C \quad (\operatorname{Re} \lambda \geq 0).$$

- Use Plancherel's Theorem to show

$$\|T(t)(-A)^{-\alpha}\| \leq Ct^{-1}.$$

- Use semigroup property and interpolation (moment inequality) to show

$$\|T(t)(-A)^{-1}\| \leq Ct^{-1/\alpha}.$$

Outline of proof: Hilbert space, regularly varying case

Hilbert space, $M(s) \sim \frac{s^\alpha}{\ell(s)}$, ℓ increasing.

- Find a complete Bernstein function f_ℓ , as large as possible, such that

$$\|(\lambda - A)^{-1}A^{-(\alpha-1)}f_\ell(-A^{-1})\| \leq C \quad (\operatorname{Re} \lambda \geq 0).$$

For example, $f_{\log}(-A^{-1}) = -A^{-1} \log(I - A)$.

- Use Plancherel's Theorem to show

$$\|T(t)A^{-(\alpha-1)}f_\ell(-A^{-1})\| \leq Ct^{-1}.$$

- Use an interpolation inequality for complete Bernstein functions of semigroup generators, together with an Abelian/Tauberian theorem for Stieltjes transforms (Karamata, 1930s), to remove the log term in the M_{\log} result.

Example of L^2 -version

X Hilbert space, $T(t)$ contractions

Assume that $D(A) = D(A^*)$. Consider $-(A + A^*)$, symmetric, non-negative.

Let S be any non-negative, self-adjoint extension of $-(A + A^*)$,
 $B = S^{1/2}$

Theorem

Assume in addition that $\sigma(A) \cap i\mathbb{R}$ is empty, and $\|R(is, A)\| \leq M(|s|)$. Then, for all $x \in X$,

$$\int_0^\infty M_{\log}^{-1}(kt)^2 \|BT(t)A^{-1}x\|^2 dt < \infty.$$

The M_{\log} -theorem, or its improvements on Hilbert space, can be applied to damped wave equations whenever the resolvent of A can be estimated. In particular,

$$\|(is - A)^{-1}\| = O(|s|^\alpha) \iff E(u, t) = O(t^{1/\alpha}).$$

The conditions of the earlier L^2 -result hold.

Corollary

Assume that the damped wave equation satisfies

$$E(u, t) \leq r(t)^2 E(u, 0)$$

for all classical solutions u , and some decreasing function $r(t)$.

Then

$$\int_0^\infty \left| \frac{d}{dt} E(u, t) \right| M_{\log}^{-1}(kt)^2 dt < \infty$$

where $M(s) = r^{-1}(c/s)$, for some $c, k > 0$.

Let \mathcal{M} be a (compact) Riemannian manifold (without boundary), and $\{\varphi_t : t \in \mathbb{R}\}$ be a smooth flow on \mathcal{M} . So-called *Anosov flows* have hyperbolic behaviour on the tangent spaces of \mathcal{M} , chaotic behaviour on \mathcal{M} , and mixing properties, i.e. for smooth f, g ,

$$\int_{\mathcal{M}} f \cdot (g \circ \varphi_t) \rightarrow \int_{\mathcal{M}} f \int_{\mathcal{M}} g \quad (t \rightarrow \infty).$$

What is the rate of convergence?

Ruelle, Pollicott, Chernoff, Dolgopyat, Liverani, Tsujii, Butterley, Faure, Sjöstrand, Dyatlov, Zworski,.....

$$C^\infty(\mathcal{M}) \subset X \subset C(\mathcal{M})$$
$$T(t)f = f \circ \varphi_t, \quad \text{generator } A$$

Ingredients:

1. Quasi-compactness argument for spectral gap
2. Resolvent estimate for large $|s|$

$$\|(\alpha + is - A)^{-\gamma \log |s|}\| \leq C(\lambda + \alpha)^{-\gamma \log |s|}. \quad (\text{Dol})$$

(fixed $\alpha, \gamma, \lambda, C > 0$ with $\gamma(\lambda + \alpha) < 1$)

(Dol) implies

$$\|(is - A)^{-1}\| = O(\log |s|)$$

and a polynomial bound for $-\lambda/2 < \text{Re } z < 0$. Exponential rate of convergence follows from (an easy case of) the M_{\log} -theorem.